

Stochastic Simulation

- **Idea:** probabilities \leftrightarrow samples
- Get probabilities from samples:

X	<i>count</i>
x_1	n_1
\vdots	\vdots
x_k	n_k
<i>total</i>	m

 \leftrightarrow

X	<i>probability</i>
x_1	n_1/m
\vdots	\vdots
x_k	n_k/m

Stochastic Simulation

- **Idea:** probabilities \leftrightarrow samples
- Get probabilities from samples:

X	<i>count</i>
x_1	n_1
\vdots	\vdots
x_k	n_k
<i>total</i>	m

 \leftrightarrow

X	<i>probability</i>
x_1	n_1/m
\vdots	\vdots
x_k	n_k/m

- If we could sample from a variable's (posterior) probability, we could estimate its (posterior) probability.

Generating samples from a distribution

For a variable X with a discrete domain or a (one-dimensional) real domain:

- Totally order the values of the domain of X .

Generating samples from a distribution

For a variable X with a discrete domain or a (one-dimensional) real domain:

- Totally order the values of the domain of X .
- Generate the cumulative probability distribution:
 $f(x) = P(X \leq x)$.

Generating samples from a distribution

For a variable X with a discrete domain or a (one-dimensional) real domain:



- Totally order the values of the domain of X .
- Generate the cumulative probability distribution:
 $f(x) = P(X \leq x)$.
- Select a value y uniformly in the range $[0, 1]$.

Generating samples from a distribution

For a variable X with a discrete domain or a (one-dimensional) real domain:

- Totally order the values of the domain of X .
- Generate the cumulative probability distribution:
 $f(x) = P(X \leq x)$.
- Select a value y uniformly in the range $[0, 1]$.
- Select the x such that $f(x) = y$.

Cumulative Distribution



Hoeffding's inequality

Theorem (Hoeffding): Suppose p is the true probability, and s is the sample average from n independent samples; then

$$P(|s - p| > \epsilon) \leq 2e^{-2n\epsilon^2}.$$

Guarantees a **probably approximately correct** estimate of probability.

Hoeffding's inequality

Theorem (Hoeffding): Suppose p is the true probability, and s is the sample average from n independent samples; then

$$P(|s - p| > \epsilon) \leq 2e^{-2n\epsilon^2}.$$

Guarantees a **probably approximately correct** estimate of probability.

If you are willing to have **an error greater than ϵ in less than δ of the cases**, solve $2e^{-2n\epsilon^2} < \delta$ for n , which gives

Hoeffding's inequality

Theorem (Hoeffding): Suppose p is the true probability, and s is the sample average from n independent samples; then

$$P(|s - p| > \epsilon) \leq 2e^{-2n\epsilon^2}.$$

Guarantees a **probably approximately correct** estimate of probability.

If you are willing to have **an error greater than ϵ in less than δ of the cases**, solve $2e^{-2n\epsilon^2} < \delta$ for n , which gives

$$n > \frac{-\ln \frac{\delta}{2}}{2\epsilon^2}.$$

Hoeffding's inequality

Theorem (Hoeffding): Suppose p is the true probability, and s is the sample average from n independent samples; then

$$P(|s - p| > \epsilon) \leq 2e^{-2n\epsilon^2}.$$

Guarantees a **probably approximately correct** estimate of probability.

If you are willing to have **an error greater than ϵ in less than δ of the cases**, solve $2e^{-2n\epsilon^2} < \delta$ for n , which gives

$$n > \frac{-\ln \frac{\delta}{2}}{2\epsilon^2}.$$

ϵ	δ	n
0.1	0.05	185
0.01	0.05	18,445
0.1	0.01	265

Forward sampling in a belief network

- Sample the variables one at a time; sample parents of X before sampling X .
- Given values for the parents of X , sample from the probability of X given its parents.

Rejection Sampling

- To estimate a posterior probability given evidence $Y_1 = v_1 \wedge \dots \wedge Y_j = v_j$:
- Reject any sample that assigns Y_i to a value other than v_i .

Rejection Sampling

- To estimate a posterior probability given evidence $Y_1 = v_1 \wedge \dots \wedge Y_j = v_j$:
- Reject any sample that assigns Y_i to a value other than v_i .
- The non-rejected samples are distributed according to the posterior probability:


$$P(\alpha \mid \text{evidence}) \approx \frac{\sum_{\alpha \text{ is true in sample}} 1}{\sum_{\text{sample}} 1}$$

where we consider only samples consistent with evidence.

Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$


	Ta	Fi	Al	Sm	Le	Re
s_1	false	true	false	true	false	false



Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$


	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	X



Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$


	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	X
s_2	false	true	true	true	true	true	



Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$


	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	✗
s_2	false	true	true	true	true	true	✓



Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$


	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	✗
s_2	false	true	true	true	true	true	✓
s_3	true	false	true	false			



Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$


	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	X
s_2	false	true	true	true	true	true	✓
s_3	true	false	true	false	—	—	X



Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$


	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	X
s_2	false	true	true	true	true	true	✓
s_3	true	false	true	false	—	—	X
s_4	true	true	true	true	true	true	



Rejection Sampling Example: $P(ta \mid sm, re)$


Observe $Sm = true, Re = true$

	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	✗
s_2	false	true	true	true	true	true	✓
s_3	true	false	true	false	—	—	✗
s_4	true	true	true	true	true	true	✓



Rejection Sampling Example: $P(ta \mid sm, re)$


Observe $Sm = true, Re = true$



	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	✗
s_2	false	true	true	true	true	true	✓
s_3	true	false	true	false	—	—	✗
s_4	true	true	true	true	true	true	✓
...							
s_{1000}	false	false	false	false			

Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$



	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	✗
s_2	false	true	true	true	true	true	✓
s_3	true	false	true	false	—	—	✗
s_4	true	true	true	true	true	true	✓
...							
s_{1000}	false	false	false	false	—	—	✗

$$P(sm) = 0.02$$

$$P(re \mid sm) = 0.32$$


There are 1000 samples.

How many are rejected?

How many are used?

Rejection Sampling Example: $P(ta \mid sm, re)$

Observe $Sm = true, Re = true$



	Ta	Fi	Al	Sm	Le	Re	
s_1	false	true	false	true	false	false	✗
s_2	false	true	true	true	true	true	✓
s_3	true	false	true	false	—	—	✗
s_4	true	true	true	true	true	true	✓
...							
s_{1000}	false	false	false	false	—	—	✗

$$P(sm) = 0.02$$

$$P(re \mid sm) = 0.32$$

There are 1000 samples.

How many are rejected?

How many are used?

Doesn't work well when evidence is unlikely.

Importance Sampling

- Samples have weights: a real number associated with each sample that takes the evidence into account.

Importance Sampling

- Samples have weights: a real number associated with each sample that takes the evidence into account.
- Probability of a proposition is weighted average of samples:

$$P(\alpha \mid \textit{evidence}) \approx \frac{\sum_{\textit{sample}: \alpha \textit{ is true in sample}} \textit{weight}(\textit{sample})}{\sum_{\textit{sample}} \textit{weight}(\textit{sample})}$$

Importance Sampling

- Samples have weights: a real number associated with each sample that takes the evidence into account.
- Probability of a proposition is weighted average of samples:

$$P(\alpha \mid \text{evidence}) \approx \frac{\sum_{\text{sample: } \alpha \text{ is true in sample}} \text{weight}(\text{sample})}{\sum_{\text{sample}} \text{weight}(\text{sample})}$$

- Mix exact inference with sampling: don't sample all of the variables, but weight each sample according to $P(\text{evidence} \mid \text{sample})$.

Importance Sampling (Likelihood Weighting)

```
procedure likelihood_weighting( $B_n, e, H, n$ ):  
  # Approximate  $P(H \mid e)$  in belief network  $B_n$  using  $n$  samples.  
  #  $H$  has some real domain (e.g.,  $\{0, 1\}$ )  
   $mass := 0$            # mass of all samples  
   $hmass := 0$         # weighted sum of value of  $H$   
  repeat  $n$  times:  
     $weight := 1$      # weight of current sample  
    for each variable  $X_i$  in order:  
      if  $X_i = o_i$  is observed  
         $weight := weight \times P(X_i = o_i \mid parents(X_i))$   
      else assign  $X_i$  a random sample of  $P(X_i \mid parents(X_i))$   
     $mass := mass + weight$   
     $hmass := hmass + weight * (\text{value of } H \text{ in current assignment})$   
  return  $hmass/mass$ 
```

Importance Sampling (Likelihood Weighting)

```
procedure likelihood_weighting( $Bn, e, H, n$ ):  
  # Approximate  $P(H \mid e)$  in belief network  $Bn$  using  $n$  samples.  
  #  $H$  has some real domain (e.g.,  $\{0, 1\}$ )  
   $mass := 0$            # mass of all samples  
   $hmass := 0$         # weighted sum of value of  $H$   
  repeat  $n$  times:  
     $weight := 1$       # weight of current sample  
    for each variable  $X_i$  in order:  
      if  $X_i = o_i$  is observed  
         $weight := weight \times P(X_i = o_i \mid parents(X_i))$   
      else assign  $X_i$  a random sample of  $P(X_i \mid parents(X_i))$   
     $mass := mass + weight$   
     $hmass := hmass + weight * (\text{value of } H \text{ in current assignment})$   
  return  $hmass/mass$ 
```

Importance Sampling (Likelihood Weighting)

```
procedure likelihood_weighting( $Bn, e, H, n$ ):  
  # Approximate  $P(H \mid e)$  in belief network  $Bn$  using  $n$  samples.  
  #  $H$  has some real domain (e.g.,  $\{0, 1\}$ )  
   $mass := 0$            # mass of all samples  
   $hmass := 0$         # weighted sum of value of  $H$   
  repeat  $n$  times:  
     $weight := 1$       # weight of current sample  
    for each variable  $X_i$  in order:  
      if  $X_i = o_i$  is observed  
         $weight := weight \times P(X_i = o_i \mid parents(X_i))$   
      else assign  $X_i$  a random sample of  $P(X_i \mid parents(X_i))$   
     $mass := mass + weight$   
     $hmass := hmass + weight * (\text{value of } H \text{ in current assignment})$   
  return  $hmass/mass$ 
```


Importance Sampling (Likelihood Weighting)

```
procedure likelihood_weighting( $Bn, e, H, n$ ):  
  # Approximate  $P(H \mid e)$  in belief network  $Bn$  using  $n$  samples.  
  #  $H$  has some real domain (e.g.,  $\{0, 1\}$ )  
   $mass := 0$            # mass of all samples  
   $hmass := 0$         # weighted sum of value of  $H$   
  repeat  $n$  times:  
     $weight := 1$       # weight of current sample  
    for each variable  $X_i$  in order:  
      if  $X_i = o_i$  is observed  
         $weight := weight \times P(X_i = o_i \mid parents(X_i))$   
      else assign  $X_i$  a random sample of  $P(X_i \mid parents(X_i))$   
     $mass := mass + weight$   
     $hmass := hmass + weight * (\text{value of } H \text{ in current assignment})$   
  return  $hmass/mass$ 
```


Importance Sampling (Likelihood Weighting)

```
procedure likelihood_weighting( $Bn, e, H, n$ ):  
  # Approximate  $P(H \mid e)$  in belief network  $Bn$  using  $n$  samples.  
  #  $H$  has some real domain (e.g.,  $\{0, 1\}$ )  
   $mass := 0$            # mass of all samples  
   $hmass := 0$         # weighted sum of value of  $H$   
  repeat  $n$  times:  
     $weight := 1$       # weight of current sample  
    for each variable  $X_i$  in order:  
      if  $X_i = o_i$  is observed  
         $weight := weight \times P(X_i = o_i \mid parents(X_i))$   
      else assign  $X_i$  a random sample of  $P(X_i \mid parents(X_i))$   
     $mass := mass + weight$   
     $hmass := hmass + weight * (\text{value of } H \text{ in current assignment})$   
  return  $hmass/mass$ 
```

Importance Sampling Example: $P(ta \mid sm, re)$



	Ta	Fi	Al	Le	Weight
s_1	true	false	true	false	
s_2	false	true	false	false	
s_3	false	true	true	true	
s_4	true	true	true	true	
...					
s_{1000}	false	false	true	true	


$$P(sm \mid fi) = 0.9$$

$$P(sm \mid \neg fi) = 0.01$$

$$P(re \mid le) = 0.75$$

$$P(re \mid \neg le) = 0.01$$

Importance Sampling Example: $P(ta \mid sm, re)$



	Ta	Fi	Al	Le	Weight
s_1	true	false	true	false	0.01×0.01
s_2	false	true	false	false	
s_3	false	true	true	true	
s_4	true	true	true	true	
...					
s_{1000}	false	false	true	true	


$$P(sm \mid fi) = 0.9$$

$$P(sm \mid \neg fi) = 0.01$$

$$P(re \mid le) = 0.75$$

$$P(re \mid \neg le) = 0.01$$

Importance Sampling Example: $P(ta \mid sm, re)$



	Ta	Fi	Al	Le	Weight
s_1	true	false	true	false	0.01×0.01
s_2	false	true	false	false	0.9×0.01
s_3	false	true	true	true	0.9×0.75
s_4	true	true	true	true	0.9×0.75
...					
s_{1000}	false	false	true	true	0.01×0.75


$$P(sm \mid fi) = 0.9$$

$$P(sm \mid \neg fi) = 0.01$$

$$P(re \mid le) = 0.75$$

$$P(re \mid \neg le) = 0.01$$

Importance Sampling Example: $P(le \mid sm, ta, \neg re)$



$$P(ta) = 0.02$$

$$P(fi) = 0.01$$

$$P(al \mid fi \wedge ta) = 0.5$$

$$P(al \mid fi \wedge \neg ta) = 0.99$$

$$P(al \mid \neg fi \wedge ta) = 0.85$$

$$P(al \mid \neg fi \wedge \neg ta) = 0.0001$$

$$P(sm \mid fi) = 0.9$$

$$P(sm \mid \neg fi) = 0.01$$

$$P(le \mid al) = 0.88$$

$$P(le \mid \neg al) = 0.001$$

$$P(re \mid le) = 0.75$$

$$P(re \mid \neg le) = 0.01$$

Computing Expectations & Proposal Distributions

Expected value of f with respect to distribution P :

$$\mathbb{E}_P(f) = \sum_w f(w) * P(w)$$

Computing Expectations & Proposal Distributions

Expected value of f with respect to distribution P :

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w) \\ &\approx \frac{1}{n} \sum_s f(s)\end{aligned}$$

s is sampled with probability P . There are n samples.
(Expectation of variable with domain $\{0, 1\}$ is its probability.)

Computing Expectations & Proposal Distributions

Expected value of f with respect to distribution P :

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w) \\ &\approx \frac{1}{n} \sum_s f(s)\end{aligned}$$

s is sampled with probability P . There are n samples.

(Expectation of variable with domain $\{0, 1\}$ is its probability.)

$$\mathbb{E}_P(f) = \sum_w f(w) * P(w)/Q(w) * Q(w)$$

Computing Expectations & Proposal Distributions

Expected value of f with respect to distribution P :

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w) \\ &\approx \frac{1}{n} \sum_s f(s)\end{aligned}$$

s is sampled with probability P . There are n samples.
(Expectation of variable with domain $\{0, 1\}$ is its probability.)

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w)/Q(w) * Q(w) \\ &\approx \frac{1}{n} \sum_s f(s) * P(s)/Q(s)\end{aligned}$$

s is selected according to the distribution Q .

Computing Expectations & Proposal Distributions

Expected value of f with respect to distribution P :

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w) \\ &\approx \frac{1}{n} \sum_s f(s)\end{aligned}$$

s is sampled with probability P . There are n samples.
(Expectation of variable with domain $\{0, 1\}$ is its probability.)

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w)/Q(w) * Q(w) \\ &\approx \frac{1}{n} \sum_s f(s) * P(s)/Q(s)\end{aligned}$$

s is selected according to the distribution Q .
The distribution Q is called a **proposal distribution**.

Computing Expectations & Proposal Distributions

Expected value of f with respect to distribution P :

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w) \\ &\approx \frac{1}{n} \sum_s f(s)\end{aligned}$$

s is sampled with probability P . There are n samples.
(Expectation of variable with domain $\{0, 1\}$ is its probability.)

$$\begin{aligned}\mathbb{E}_P(f) &= \sum_w f(w) * P(w)/Q(w) * Q(w) \\ &\approx \frac{1}{n} \sum_s f(s) * P(s)/Q(s)\end{aligned}$$

s is selected according to the distribution Q .

The distribution Q is called a **proposal distribution**.

$P(c) > 0$ then $Q(c) > 0$.

Try to make Q so the weights end up far from zero.

Particle Filtering

Importance sampling can be seen as:

for each particle:

for each variable:

sample / absorb evidence / update query

where **particle** is one of the samples.

Particle Filtering

Importance sampling can be seen as:

for each particle:

for each variable:

sample / absorb evidence / update query

where **particle** is one of the samples.

Instead we could do:

for each variable:

for each particle:

sample / absorb evidence / update query

Particle Filtering

Importance sampling can be seen as:

for each particle:

for each variable:

sample / absorb evidence / update query

where **particle** is one of the samples.

Instead we could do:

for each variable:

for each particle:

sample / absorb evidence / update query

Why?

Particle Filtering

Importance sampling can be seen as:

for each particle:

for each variable:

sample / absorb evidence / update query

where **particle** is one of the samples.

Instead we could do:

for each variable:

for each particle:

sample / absorb evidence / update query

Why?

- It works with infinitely many variables (e.g., HMM)

Particle Filtering

Importance sampling can be seen as:

for each particle:

for each variable:

sample / absorb evidence / update query

where **particle** is one of the samples.

Instead we could do:

for each variable:

for each particle:

sample / absorb evidence / update query

Why?

- It works with infinitely many variables (e.g., HMM)
- We can have a new operation of resampling

Particle Filtering for HMMs

- Start with random chosen particles (say 1000)
- Sample initial states in proportion to their probability.

Particle Filtering for HMMs

- Start with random chosen particles (say 1000)
- Sample initial states in proportion to their probability.
- Repeat (as each observation arrives):
 - ▶ **Absorb evidence**: weight each particle by the probability of the evidence observation given the state of the particle.

Particle Filtering for HMMs

- Start with random chosen particles (say 1000)
- Sample initial states in proportion to their probability.
- Repeat (as each observation arrives):
 - ▶ **Absorb evidence**: weight each particle by the probability of the evidence observation given the state of the particle.
 - ▶ **Resample**: select each particle at random, in proportion to the weight of the particle.
Some particles may be duplicated, some may be removed. All new particles have same weight.

Particle Filtering for HMMs


- Start with random chosen particles (say 1000)
- Sample initial states in proportion to their probability.
- Repeat (as each observation arrives):
 - ▶ **Absorb evidence**: weight each particle by the probability of the evidence observation given the state of the particle.
 - ▶ **Resample**: select each particle at random, in proportion to the weight of the particle.
Some particles may be duplicated, some may be removed. All new particles have same weight.
 - ▶ **Transition**: sample the next state for each particle according to the transition probabilities.

Particle Filtering for HMMs

- Start with random chosen particles (say 1000)
- Sample initial states in proportion to their probability.
- Repeat (as each observation arrives):
 - ▶ **Absorb evidence**: weight each particle by the probability of the evidence observation given the state of the particle.
 - ▶ **Resample**: select each particle at random, in proportion to the weight of the particle.
Some particles may be duplicated, some may be removed. All new particles have same weight.
 - ▶ **Transition**: sample the next state for each particle according to the transition probabilities.

To answer a query about the current state, use the set of particles as data.

Example: Localization



Loc consists of (x, y, θ) – position and orientation
 $k = 24$ sonar sensors (all very noisy)

Markov Chain Monte Carlo

To sample from a distribution P :

- Create (ergodic and aperiodic) Markov chain with P as equilibrium distribution.
Let $T(S_{i+1} | S_i)$ be the transition probability.
- Given state s , sample state s' from $T(S | s)$

Markov Chain Monte Carlo

To sample from a distribution P :

- Create (ergodic and aperiodic) Markov chain with P as equilibrium distribution.

Let $T(S_{i+1} | S_i)$ be the transition probability.

- Given state s , sample state s' from $T(S | s)$
- After a while, the states sampled will be distributed according to P .

Markov Chain Monte Carlo

To sample from a distribution P :

- Create (ergodic and aperiodic) Markov chain with P as equilibrium distribution.
Let $T(S_{i+1} | S_i)$ be the transition probability.
- Given state s , sample state s' from $T(S | s)$
- After a while, the states sampled will be distributed according to P .
- Ignore the first samples “burn-in”
— use the remaining samples.
- Samples are not independent of each other “autocorrelation”.
Sometimes use subset (e.g., 1/1000) of them “thinning”


To sample from a distribution P :

- Create (ergodic and aperiodic) Markov chain with P as equilibrium distribution.
Let $T(S_{i+1} | S_i)$ be the transition probability.
- Given state s , sample state s' from $T(S | s)$
- After a while, the states sampled will be distributed according to P .
- Ignore the first samples “burn-in”
— use the remaining samples.
- Samples are not independent of each other “autocorrelation”.
Sometimes use subset (e.g., 1/1000) of them “thinning”
- **Gibbs sampler**: sample each non-observed variable from the distribution of the variable given the current (or observed) value of the variables in its Markov blanket.

Gibbs Sampling Example: $P(ta \mid sm, re)$

	Ta	Fi	Al	Le
s_1	true	false	false	true


Select Le .




Gibbs Sampling Example: $P(ta \mid sm, re)$

	Ta	Fi	Al	Le
s_1	true	false	false	true

Select Le . Sample from $P(Le \mid \neg al \wedge re)$




Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select Le . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false

Gibbs Sampling Example: $P(ta \mid sm, re)$



	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> .				

Gibbs Sampling Example: $P(ta \mid sm, re)$



	Ta	Fi	Al	Le
--	----	----	----	----


s_1	true	false	false	true
-------	------	-------	-------	------

Select *Le*. Sample from $P(Le \mid \neg al \wedge re)$

s_2	true	false	false	false
-------	------	-------	-------	-------


Select *Fi*. Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false
Select <i>Al</i> .				

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false
Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$				

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false
Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$				
s_4	true	true	false	false

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false
Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$				
s_4	true	true	false	false
Select <i>Le</i> .				

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false
Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$				
s_4	true	true	false	false
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false
Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$				
s_4	true	true	false	false
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_5	true	true	false	true

Gibbs Sampling Example: $P(ta \mid sm, re)$




	Ta	Fi	Al	Le
s_1	true	false	false	true
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_2	true	false	false	false
Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$				
s_3	true	true	false	false
Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$				
s_4	true	true	false	false
Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$				
s_5	true	true	false	true
Select <i>Ta</i> .				

Gibbs Sampling Example: $P(ta \mid sm, re)$



	Ta	Fi	Al	Le
s_1	true	false	false	true
	Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$			
s_2	true	false	false	false
	Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$			
s_3	true	true	false	false
	Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$			
s_4	true	true	false	false
	Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$			
s_5	true	true	false	true
	Select <i>Ta</i> . Sample from $P(Ta \mid \neg al \wedge fi)$			

Gibbs Sampling Example: $P(ta \mid sm, re)$



	Ta	Fi	Al	Le
s_1	true	false	false	true
	Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$			
s_2	true	false	false	false
	Select <i>Fi</i> . Sample from $P(Fi \mid ta \wedge \neg al \wedge sm)$			
s_3	true	true	false	false
	Select <i>Al</i> . Sample from $P(Al \mid ta \wedge fi \wedge \neg le)$			
s_4	true	true	false	false
	Select <i>Le</i> . Sample from $P(Le \mid \neg al \wedge re)$			
s_5	true	true	false	true
	Select <i>Ta</i> . Sample from $P(Ta \mid \neg al \wedge fi)$			
s_6	true	true	false	true
...				

